

# Mirage unification at TeV scale and natural electroweak symmetry breaking in minimal supersymmetry

Ken-ichi Okumura

*Department of Physics, Kyushu University, Fukuoka 812-8581, Japan*

**Abstract.** We propose a new minimal supersymmetric scenario with virtually no fine-tuning in the electroweak symmetry breaking. It favors light supersymmetric spectrum below a few TeV and predicts definite relations among stop and gaugino masses and LSP higgsino with  $\mu \lesssim 200$  GeV, which are within the reach of coming LHC experiment <sup>1</sup>.

**Keywords:** supersymmetry, electroweak symmetry breaking, mirage mediation

Supersymmetry (SUSY) is a leading candidate for physics beyond the standard model (SM) which solves the enigmatic gauge hierarchy problem. The minimal extension leads to various non-trivial predictions: gauge coupling unification, radiative electroweak (EW) symmetry breaking, the lightest SUSY particle (LSP) protected by R-parity (a cold dark matter candidate) and a light Higgs boson compatible with the EW precision tests. All of them are direct consequences of a single spell, 'minimal supersymmetry'. This is a remarkable success, considering various difficulties we often encounter in alternative approaches.

However, the LEP II experiment observed neither a Higgs boson nor a SUSY particle and has built up an uncomfortable tension in the EW symmetry breaking of the minimal supersymmetric standard model (MSSM). At tree level, the lightest Higgs boson in the MSSM is lighter than  $M_Z$ . Thus, the LEP II SM Higgs mass bound  $m_{h^0}^2 > 114.4$  GeV requires a substantial radiative correction, which raises the upper bound approximately,

$$m_{h^0}^2 \lesssim M_Z^2 + (3g^2 m_t^4 / 8\pi^2 M_W^2) \ln(m_{\tilde{t}}^2 / m_t^2), \quad (1)$$

neglecting the trilinear scalar coupling,  $A_t$ . Then the SM Higgs mass bound is translated into  $m_{\tilde{t}} \gtrsim 500$  GeV. This bound for the stop mass is closely related to that of the up-type Higgs,  $H_u$  through the renormalization group (RG) running. If the boundary value of  $m_{H_u}^2$  is given at the cut-off scale  $\Lambda$ , a dominant correction below it is estimated as,

$$\Delta m_{H_u}^2 \sim - (3/4\pi^2) y_t^2 m_{\tilde{t}}^2 \ln(\Lambda / m_{\tilde{t}}). \quad (2)$$

Then  $\Lambda \approx M_{GUT}$  leads to  $\Delta m_{H_u}^2 \sim -2m_{\tilde{t}}^2$ , which suggests  $m_{H_u}^2 \sim \mathcal{O}(m_{\tilde{t}}^2)$  up to a 'conspiracy' between  $m_{H_u}^2(\Lambda)$  and the RG running. While the EW symmetry breaking requires,

$$M_Z^2/2 = (m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta) / (\tan^2 \beta - 1) - |\mu|^2 \approx -m_{H_u}^2 - |\mu|^2. \quad (3)$$

---

<sup>1</sup> This talk is based on the collaboration in [1]

Unless  $m_{H_u}^2 \sim \mu^2 \sim M_Z^2$ , we need a fine-tuning between  $m_{H_u}^2$  and  $\mu^2$ , despite their origins are totally different from each other. The degree of fine-tuning is measured by,

$$\Delta(\mu^2)^{-1} \equiv (\partial \ln M_Z^2 / \partial \ln \mu^2)^{-1} = M_Z^2 / 2|\mu|^2. \quad (4)$$

This implies  $m_{H_u} \sim m_{\tilde{t}} \gtrsim 500$  GeV results in a fine-tuning worse than a few % level.

Various solutions have been proposed for this 'little SUSY hierarchy problem' between the EW scale and the SUSY (stop) scale. They are classified in two categories. The first solution raises the tree level Higgs mass well above  $M_Z$  by enhancing the Higgs quartic coupling, which is solely determined by  $SU(2)_L \times U(1)_Y$  D-term in the MSSM. The second solution explores a way to realize the little hierarchy by introducing a new symmetry around the EW scale or lowering the cut-off  $\Lambda$  close to the EW scale. In both cases, new fields and thresholds are required. This spoils the original simplicity of the minimal scenario, in particular, the prediction of the gauge coupling unification.

In this paper, we pursue an alternative direction, preparing the conspiracy between  $m_{H_u}^2(M_{GUT})$  and the RG running below it, in order to realize the EW scale little hierarchy [1, 2]. This looks hard to achieve but is actually possible in the mirage mediation scenario of SUSY breaking [3, 4, 5], typically realized in the KKLT-type string modulus stabilization[6]. The system is described by the following effective supergravity action,

$$\int d^4x \left[ \int d^4\theta (-3)CC^* \exp(-\mathcal{K}/3) + \left\{ \int d^2\theta ((f_a/4)W^a W^a + C^3 \mathcal{W}) + \text{h.c.} \right\} \right. \\ \left. + \int d^4\theta C^2 C^{*2} \mathcal{P}_{\text{lift}} \theta^2 \bar{\theta}^2 \right] \quad (5)$$

where  $C = C_0 + F^C \theta^2$  denotes the chiral compensator superfield. Kähler potential  $\mathcal{K}$ , gauge kinetic function  $f_a$ , superpotential  $\mathcal{W}$  and uplifting function  $\mathcal{P}_{\text{lift}}$  (represents the effect of SUSY breaking brane [6, 3]) are given by,

$$\begin{aligned} \mathcal{K} &= \mathcal{K}_0(T + T^*) + Z_i(T + T^*)\Phi_i^* \Phi_i, & f_a &= T, \\ \mathcal{W} &= w_0 - A e^{-aT} + \lambda_{ijk} \Phi_i \Phi_j \Phi_k, & \mathcal{P}_{\text{lift}} &= \mathcal{P}_{\text{lift}}(T + T^*), \end{aligned} \quad (6)$$

where  $T$  denotes the gauge coupling modulus and  $\Phi_i$  stands for chiral matter superfields. In  $\mathcal{P}_{\text{lift}} = 0$  limit, the modulus is stabilized at SUSY AdS vacuum with  $aT \approx \ln(A/w_0) \approx \ln(M_{Pl}/m_{3/2})$ , where  $F^T \equiv -e^{\mathcal{K}/2} K^{TT^*} D_T^* \mathcal{W}^* = 0$  and the potential minimum is given by  $\langle V \rangle = -3|m_{3/2}|^2$ . This vacuum energy is fine-tuned to  $\langle V \rangle \approx 0^+$  by  $\mathcal{P}_{\text{lift}}$ . This uplifting process generates finite  $M_0 \equiv F^T/(T + T^*)$  due to the shift of modulus vev [3]. We introduce a parameter,  $\alpha$  indicating the relative size of  $m_{3/2}$  to  $M_0$ ,

$$\alpha \equiv m_{3/2}/M_0 \ln(M_{Pl}/m_{3/2}) \approx \{1 + (3/2)(\partial_T \ln \mathcal{P}_{\text{lift}} / \partial_T \mathcal{K}_0)\}^{-1}. \quad (7)$$

This is typically  $\mathcal{O}(1)$  and modulus mediated SUSY breaking in the visible sector is comparable to the anomaly mediated one [3]. Note that the relative phase is dynamically eliminated [7, 5]. The soft SUSY breaking terms of the visible fields are summarized as,

$$\mathcal{L} = -(1/2) M_a \bar{\lambda}^a \lambda^a - m_i^2 |\phi_i|^2 - \{(1/6) A_{ijk} y_{ijk} \phi_i \phi_j \phi_k + \text{h.c.}\}. \quad (8)$$

where  $\phi_i$  denotes a scalar component of  $\Phi_i$ , and  $\lambda^a$  represents gaugino. The canonical Yukawa coupling  $y_{ijk}$  is defined as,  $y_{ijk} = \lambda_{ijk}/\sqrt{e^{-\mathcal{K}_0} Z_i Z_j Z_k}$ . These terms are derived at the unification scale using standard supergravity formula,

$$\begin{aligned} M_a &= M_0 + \frac{b_a g_a^2}{16\pi^2} m_{3/2}, \quad A_{ijk} = (a_i + a_j + a_k) M_0 - \frac{1}{16\pi^2} (\gamma_i + \gamma_j + \gamma_k) m_{3/2}, \\ m_i^2 &= c_i M_0^2 - \frac{m_{3/2} M_0}{8\pi^2} \left( 2C_a^a(\phi_i) g_a^2 - \sum_{jk} \frac{1}{2} |y_{ijk}|^2 (a_i + a_j + a_k) \right) - \frac{1}{32\pi^2} \frac{d\gamma_i}{d\ln\mu} |m_{3/2}|^2, \\ \gamma_i &= (8\pi^2) \frac{d\ln Z_i}{d\ln\mu}, \quad a_i = 2\text{Re}(T) \partial_T \ln(e^{-\mathcal{K}_0/3} Z_i), \quad c_i = -4\text{Re}(T)^2 \partial_T \partial_{T^*} \ln(e^{-\mathcal{K}_0/3} Z_i), \end{aligned} \quad (9)$$


where  $a_i$  and  $c_i$  are typically non-negative rational numbers.

Intriguing feature of this mirage mediation is that the anomaly mediated contribution ( $\propto m_{3/2}, m_{3/2}^2$ ) effectively shifts the modulus mediation scale from  $M_{GUT}$  to the *mirage messenger scale*,  $M_{\text{mir}} \equiv M_{GUT}/(M_{\text{Pl}}/m_{3/2})^{\alpha/2}$  at one-loop level [4]. This holds for all the soft terms if the Yukawa coupling is negligible or only allowed for the combination of fields which satisfies  $a_i + a_j + a_k = c_i + c_j + c_k = 1$ . It is straightforward to show  $M_a(\mu) = (g_a^2(\mu)/g_a^2(M_{\text{mir}})) M_0$  by explicit calculation. While at  $M_{GUT}$ ,  $g_a^2 = g_G^2 = \text{Re}(T)^{-1}$  and  $y_{ijk}^2 = \lambda^2/e^{-\mathcal{K}_0} Z_i Z_j Z_k \propto \text{Re}(T)^{-1}$ , if the above condition is satisfied. Hence, the anomalous dimension scales like  $\gamma_i(\text{Re}(T)) = \text{Re}(T)^{-1} \gamma_i(1)$  at one-loop level. This enables us to solve the  $T$  dependence of  $\ln Z_i$  at arbitrary scale as  $\ln Z_i(\text{Re}(T))|_{\mu=M_{GUT}} + \Delta \ln Z_i((\mu/M_{GUT})^{1/g_G^2 \text{Re}(T)})$ . Promoting  $T$  as a superfield and replacing  $M_{GUT} \rightarrow M_{GUT} \sqrt{CC^*}$ , it is straightforward to show  $\Delta \ln Z_i|_{\theta^2} = \Delta \ln Z_i|_{\theta^2 \bar{\theta}^2} = 0$  at  $M_{\text{mir}}$ , which leads to  $A_{ijk}(M_{\text{mir}}) = M_0$  and  $m_i^2(M_{\text{mir}}) = c_i M_0^2$  in case of non-Abelian gauge group. Then the soft terms having the same  $c_i$  unify at  $M_{\text{mir}}$  (*mirage unification*).

This property is exactly what we need to realize the little hierarchy between the Higgs and stop masses in the minimal SUSY. Once we have  $\alpha = 2$ ,  $a_{u_3} + a_{q_3} + a_{H_u} = 1$  and  $c_{H_u} = 0$ ,  $c_{u_3} + c_{q_3} = 1$  due to discrete nature of the underlying physics, we can realize  $M_{\text{mir}} \approx M_0$  and  $m_{H_u}^2(M_{\text{mir}}) \approx M_0^2/8\pi^2$  while  $m_{\tilde{u}_3}^2(M_{\text{mir}}) + m_{\tilde{q}_3}^2(M_{\text{mir}}) = M_0^2$ . Note that we do not add any new fields other than  $T$  and the gauge coupling constants unify at  $M_{GUT}$ .

An extra bonus of this scenario is natural realization of  $\mu$  and  $B$  of order  $M_0$ , which is non-trivial to achieve with the anomaly mediation due to an unsuppressed  $\mathcal{O}(m_{3/2})$  contribution to  $B$ . If we assume the  $\mu$  term is generated by the same non-perturbative dynamics as in the modulus stabilization, like  $\Delta \mathcal{W} = \tilde{A} e^{-aT} H_d H_u$ ,  $\mu$  is naturally  $\mathcal{O}(M_0)$  and the leading  $\mathcal{O}(m_{3/2})$  terms in  $B$  cancel at  $\alpha = 2$ , leaving  $B = \mathcal{O}(M_0)$  [4, 1].

In this scenario,  $H_d$  can belong to either the EW scale ( $c_{H_d} = 0$ ) or the SUSY scale ( $c_{H_d} \neq 0$ ) as far as  $\tan\beta$  is not extremely large. In the former case, the EW symmetry breaking condition,  $B\mu \approx (m_{H_d}^2 + m_{H_u}^2 + 2\mu^2)/\tan\beta$  requires  $B \sim M_Z/\tan\beta$ . This is beyond the control of the above mechanism generating  $B$ , although it is not the problem of the effective theory description itself. While, in the latter case,  $B \sim M_0$  and virtually no fine-tuning is required. It is known that this non-universal choice for  $c_{H_d, H_u}$  generates an effective  $U(1)$  D-term contribution in the soft masses. It is positive for  $m_{H_u}^2$ , however, sufficiently small to be driven negative by other corrections in similar size. In the left



**FIGURE 1.** RG evolution and fine-tuning of the EW symmetry breaking in the TeV mirage mediation.

panel of Fig.1, we show RG evolution of the soft masses for  $c_{q,u,d,l,e} = 1/2$ ,  $c_{H_d} = 1$ . Non-negligible bottom Yukawa coupling and the effective D-term contribution perturb the exact unification, however, a sufficient little hierarchy is realized at  $\approx M_0$ . In the right panel of Fig.1, we plot the fine-tuning parameters  $\Delta(\mu^2)^{-1}$  and  $\Delta(|B|)^{-1}$  (similar definition as in (4)) as a function of  $m_{H_u}^2 (M_0/\sqrt{2})$ . Here  $B_0$  denotes  $B(M_{GUT})$  other than the loop induced anomaly mediated contribution. Note that the EW symmetry is broken with fine-tuning better than 10 % for predicted  $m_{H_u}^2 \sim M_0^2/8\pi^2$  with  $M_0 \approx M_{1,2,3} \gtrsim 1$  TeV while the lightest Higgs mass is comfortably accommodated above the SM bound. In this scenario, LSP is always pure higgsino, typically lighter than  $\sim 200$  GeV.

## ACKNOWLEDGMENTS

This work is completed under the grant-in-aid for scientific research on priority areas (No. 441): "Progress in elementary particle physics of the 21 century through discoveries of Higgs boson and supersymmetry" (No. 16081209) from the MECSST of Japan. The numerical calculations were carried out on Altix3700 BX2 at YITP in Kyoto University.

## REFERENCES

1. K. Choi, K.S. Jeong, T. Kobayashi and K.i. Okumura, Phys. Lett. B **633**, 355 (2006) [arXiv:hep-ph/0508029]; in preparation.
2. R. Kitano and Y. Nomura, Phys. Lett. B **631**, 58 (2005) [arXiv:hep-ph/0509039]; O. Lebedev, H. P. Nilles and M. Ratz, arXiv:hep-ph/0511320; A. Pierce and J. Thaler, arXiv:hep-ph/0604192.
3. K. Choi, A. Falkowski, H. P. Nilles, M. Olechowski and S. Pokorski, JHEP **0411**, 076 (2004) [arXiv:hep-th/0411066]; K. Choi, A. Falkowski, H. P. Nilles and M. Olechowski, Nucl. Phys. B **718**, 113 (2005) [arXiv:hep-th/0503216]; O. Loaiza-Brito, J. Martin, H. P. Nilles and M. Ratz, AIP Conf. Proc. **805**, 198 (2006) [arXiv:hep-th/0509158].
4. K. Choi, K. S. Jeong and K. i. Okumura, JHEP **0509**, 039 (2005) [arXiv:hep-ph/0504037]; K. i. Okumura, arXiv:hep-ph/0509225.
5. M. Endo, M. Yamaguchi, K. Yoshioka, Phys. Rev. D **72**, 015004 (2005) [arXiv:hep-ph/0504036].
6. S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0301240].
7. K. Choi, Phys. Rev. Lett. **72**, 1592 (1994) [arXiv:hep-ph/9311352].